

AD-A121 731

CATEGORY ANALYSIS OF THE MARINE CORPS COMBAT READINESS
EVALUATION SYSTEM (U) GEORGE WASHINGTON UNIV
WASHINGTON DC INST FOR MANAGEMENT SCIE..

1/1

UNCLASSIFIED

S ZACKS ET AL. 15 OCT 81 SERIAL-T-458


F/G 15/7.

NL

END

FMHQ

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A121731

THE
GEORGE
WASHINGTON
UNIVERSITY

STUDENTS FACULTY STUDY
RESEARCH DEVELOPMENT FUTURE CAREER CREATIVITY COMMUNITY LEADERSHIP TECHNOLOGY FRONTIER DESIGN ENGINEERING APPLIED SCIENCE
GEORGE WASHINGTON UNIVERSITY

NOV 23 1982

A

8

SCHOOL OF ENGINEERING
AND APPLIED SCIENCE

THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE: ITS DISTRIBUTION IS UNLIMITED

DUPLICATE COPY



CATEGORY ANALYSIS OF THE MARINE CORPS
COMBAT READINESS EVALUATION SYSTEM (MCCRES)

by

S. Zacks, W. H. Marlow and Zeev Barzily

Serial T-450 ✓
15 October 1981

The George Washington University
School of Engineering and Applied Science
Institute for Management Science and Engineering

Program in Logistics
Contract N00014-75-C-0729
Project NR 347 020
Office of Naval Research

DTIC
SELECT
NOV 25 1982
A

This document has been approved for public
sale and release; its distribution is unlimited.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER T-450	2. GOVT ACCESSION NO. AD A/21731	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Category Analysis of the Marine Corps Combat Readiness Evaluation System (MCCRES)		5. TYPE OF REPORT & PERIOD COVERED Scientific
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) S. Zacks W. H. Marlow Zeev Barzily		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0729
9. PERFORMING ORGANIZATION NAME AND ADDRESS THE GEORGE WASHINGTON UNIVERSITY PROGRAM IN LOGISTICS WASHINGTON, DC 20052		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Code 411MA Arlington, VA 22217		12. REPORT DATE 15 October 1981
		13. NUMBER OF PAGES 24
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public sale and release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Marine Corps Readiness Multivariate Analysis Infantry Readiness Readiness		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Twenty-seven data sets (evaluations) of the MCCRES are analyzed from the point of view of the category structure introduced by Barzily. Due to the specific designs of the evaluations, the statistical data are complicated by missing observations (nonapplicable requirements). This creates a problem of analyzing dependent Bernoulli trials with different numbers of observations in different cells. A quasi-Bayesian model is introduced and estimators of the structural parameters are studied.		

DD FORM 1473
1 JAN 75EDITION OF 1 NOV 65 IS OBSOLETE
S/N 1102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

THE GEORGE WASHINGTON UNIVERSITY
School of Engineering and Applied Science
Institute for Management Science and Engineering

Program in Logistics

Abstract
of
Serial T-450
15 October 1981

CATEGORY ANALYSIS OF THE MARINE CORPS
COMBAT READINESS EVALUATION SYSTEM (MCCRES)

by

S. Zacks, W. H. Marlow and Zeev Barzily

Twenty-seven data sets (evaluations) of the MCCRES are analyzed from the point of view of the category structure introduced by Barzily. Due to the specific designs of the evaluations, the statistical data are complicated by missing observations (nonapplicable requirements). This creates a problem of analyzing dependent Bernoulli trials with different numbers of observations in different cells. A quasi-Bayesian model is introduced and estimators of the structural parameters are studied.



Research Supported by
Contract N00014-75-C-0729
Project NR 347 020
Office of Naval Research

SEARCHED	INDEXED
SERIALIZED	FILED
OCT 15 1981	
FBI - NEW YORK	

A

THE GEORGE WASHINGTON UNIVERSITY
School of Engineering and Applied Science
Institute for Management Science and Engineering
Program in Logistics

CATEGORY ANALYSIS OF THE MARINE CORPS
COMBAT READINESS EVALUATION SYSTEM (MCCRES)

by

S. Zacks, W. H. Marlow and Zeev Barzily

1. Introduction

The Marine Corps Combat Readiness Evaluation System (MCCRES) and various analyses have been discussed recently in a series of papers by Barzily, Catalogne and Marlow [2], Zacks and Marlow [7] and Barzily, Marlow, and Zacks [8]. In the present paper we conduct statistical analysis of MCCRES from the point of view of the ten categories introduced by Barzily [1]. As in the previous work we restrict our present attention to Volume II of MCCRES, which deals with infantry battalions and contains 800 different requirements. The category approach of Barzily classifies all requirements into the following ten categories.

1. REPORTING to higher levels of command
2. PREPARING for operation
3. COMMUNICATING
4. PERFORMING as Marines
5. DELIVERING supporting fire
6. PLANNING of operations
7. CONFORMING to doctrine

8. EXECUTING operations
9. PROVIDING combat service support
10. SUPERVISING required actions of individual Marines

In a given evaluation an infantry battalion is evaluated on a subset of the 800 requirements. That is, due to time, budget, and other constraints, not all 800 requirements are applicable. The battalion is given a score of "Yes" in case a requirement is fulfilled satisfactorily, and a score of "No" otherwise. The basic assumption in our category approach is that the probability of "Yes" for a given unit is the same for all the applicable requirements of the same category (see Appendix 1). These probabilities may change from one category to another. The vector of the ten probabilities, corresponding to the ten categories, is a parametric vector which characterizes the unit at that particular evaluation. Different units usually have different characterizing parametric vectors and the same unit may have different characterizing parametric vectors in different evaluations. In addition to the parametric vector of "Yes" probabilities, there are several other interesting statistical features of the data. The proportion of "Yes's" among the applicable requirements of different categories in the same evaluation are not independent. It is interesting to study the correlation structure between these categories. In the present study we present statistical models and develop procedures designed to estimate interesting model parameters. We base the analysis on the results of $N = 27$ evaluations. The statistical methods developed here have a wide range of possible applications for the analysis of categorical data in which a different number of independent Bernoulli trials are performed in each category, but the proportions of successes in different categories are dependent variables.

2. The Structure of the Data and the Statistical Model

Let N designate the number of evaluations, $N = 27$, and k designate the number of categories, $k = 10$. Let X_{ij} , $i=1, \dots, N$, $j=1, \dots, k$ designate the number of "Yes's" among the applicable requirements of the

j -th category in the i -th evaluation. Let M_{ij} , $i=1, \dots, N$, $j=1, \dots, k$ designate the corresponding number of applicable requirements. In Table 2.1 we present the values of X_{ij} and M_{ij} which have been compiled from MCCRES data files. All the analyses in the present paper are based on the data in this table.

Let \underline{X}_i ($i=1, \dots, N$) designate the vector of k X - values corresponding to evaluation i , and \underline{M}_i the vectors of corresponding M - values. The fundamental assumptions of the statistical model are:

- A.1 $\underline{X}_1, \underline{X}_2, \dots, \underline{X}_N$ are independent random vectors;
- A.2 The marginal distribution of X_{ij} is binomial,
 $B(M_{ij}, \theta_{ij})$, for all $i=1, \dots, N$, and $j=1, \dots, k$.
 θ_{ij} designates the probability of "Yes" for all
the requirements in evaluation i and category j .

In Appendix 1 we test a null hypothesis which is based on assumption A.2. This hypothesis is accepted by the data in such a way to provide substantial empirical verification.

The multivariate distributions of \underline{X}_i , $i=1, \dots, N$, are specified indirectly by considering first the following transformations. Let

$$Y_{ij} = 2 \sin^{-1} \left(\sqrt{\frac{X_{ij} + 1/2}{M_{ij} + 1}} \right), \quad \begin{matrix} i=1, \dots, N \\ j=1, \dots, k \end{matrix} \quad (2.1)$$

These transformations are known as variance stabilizing transformations for binomial random variables (see Johnson and Kotz [3; p. 65], who suggest the transformation $Y = 2 \sin^{-1} \sqrt{\hat{\theta}}$, where $\hat{\theta} = (x + 3/8)/(M + 3/4)$. The asymptotic properties of (2.1) and of this one are the same).

Table 2.1

Total number of satisfied and applicable requirements
by category and evaluation*

EVAL/CAT.	1	2	3	4	5	6	7	8	9	10
1	62	44	40	14	55	100	50	53	24	13
	67	54	40	23	69	110	63	73	32	17
2	79	70	42	18	67	133	65	97	31	17
	80	76	44	24	72	140	73	100	37	20
3	41	19	26	12	45	38	44	47	15	12
	42	26	28	21	56	45	53	54	16	14
4	60	33	25	20	53	68	49	68	18	16
	62	36	31	24	55	71	54	72	19	16
5	35	34	35	21	52	93	48	46	18	12
	52	44	37	24	54	99	54	67	21	14
6	52	29	20	23	57	60	59	62	18	18
	55	29	29	24	63	63	61	67	18	18
7	44	41	32	15	63	90	47	50	30	13
	47	46	34	21	64	93	51	55	31	17
8	29	18	28	12	48	40	34	65	7	9
	60	48	32	22	63	81	57	87	14	15
9	64	41	36	25	64	78	44	93	16	18
	71	48	37	26	64	90	57	104	18	18
10	64	48	35	19	60	99	45	67	25	17
	70	55	38	24	67	119	61	77	30	17
11	15	4	23	18	56	15	30	17	13	11
	15	4	23	18	56	15	31	17	13	12
12	12	5	14	16	35	10	16	12	8	7
	14	8	17	16	54	16	30	18	10	10
13	11	11	5	6	6	15	8	36	2	5
	11	11	6	9	6	16	10	37	2	5
14	54	31	23	17	57	55	49	62	16	13
	54	35	27	25	59	62	57	70	17	18
15	41	21	13	16	4	36	33	44	13	11
	42	23	17	23	4	36	35	47	13	16
16	49	35	29	13	59	73	59	63	18	12
	54	38	32	23	59	74	63	70	18	16
17	44	29	18	16	9	52	44	54	18	13
	47	30	21	23	9	52	45	55	19	13
18	69	40	35	26	62	76	66	87	22	14
	69	41	36	28	63	78	68	90	22	16
19	78	43	38	27	64	90	60	92	19	17
	79	51	38	27	65	98	71	108	20	17
20	48	26	25	13	58	49	38	56	16	17
	54	31	29	24	61	61	56	67	18	18
21	18	5	14	4	41	14	22	12	12	9
	24	11	21	17	56	22	36	28	14	15
22	71	35	29	21	52	85	45	85	24	19
	73	37	34	25	66	92	54	90	24	20
23	71	41	33	26	63	87	63	93	21	15
	73	41	36	28	64	88	68	97	23	16
24	47	30	28	22	55	61	52	49	13	16
	49	31	30	25	57	63	57	58	15	17
25	44	32	29	16	31	56	50	47	15	16
	48	34	32	25	61	64	58	58	16	18
26	34	29	24	21	52	48	32	42	15	14
	37	32	27	23	58	56	40	54	15	15
27	100	58	32	27	69	121	75	136	26	18
	101	58	38	29	71	122	78	138	28	19

*The upper number in each cell represents the total number of satisfied requirements; the lower number represents the total number of applicable requirements.

For large values of M_{ij} the random variables Y_{ij} are distributed approximately normally, with mean $\eta_{ij} = 2 \sin^{-1} \sqrt{\theta_{ij}}$ and variance $1/M_{ij}$. Let $\underline{Y}_i = (Y_{i1}, \dots, Y_{ik})'$ be the k -variate vector of the transformed X -values of the i -th evaluations. According to assumption A.1, $\underline{Y}_1, \dots, \underline{Y}_N$ are independent random vectors. We introduce here a third assumption concerning the multivariate distribution of \underline{Y}_i ($i=1, \dots, N$).

A.3 The random vectors \underline{Y}_i have multivariate normal distributions with means $\underline{\eta}_i = (\eta_{i1}, \dots, \eta_{ik})'$ and covariance matrices $\Phi_i = (\Phi_{i.jj'})$,

where

$$\Phi_{i.jj'} = \begin{cases} \frac{1}{M_{ij}} & , \quad \text{if } j=j' \\ \rho_{jj'} / \sqrt{M_{ij} M_{ij'}} & , \quad \text{if } j \neq j' \end{cases} \quad (2.2)$$

$\rho_{jj'}$ is the correlation between Y_{ij} and $Y_{ij'}$.

It is assumed that the correlations $\rho_{jj'}$ are the same for all evaluations. In Appendix 2 we show that $\rho_{jj'}$ are asymptotically equivalent to the correlations between the proportions of Yes's, $\hat{\theta}_{ij} = X_{ij}/M_{ij}$.

3. The Bayesian Framework

The Bayesian model assumes that the characterizing parametric vectors $\underline{\theta}_i$ of the different evaluations vary according to a (prior) distribution of all possible evaluations. This is a model of a superpopulation which makes it possible to relate the information obtained from all evaluations in a consistent manner. Following the model developed in the previous section, we present here the Bayesian model in terms of the distributions of \underline{Y}_i ($i=1, \dots, N$).

We assume that the mean vectors $\eta_{i\cdot}$ of the distributions of $Y_i (i=1, \dots, N)$ have a prior normal distribution with mean vector $\bar{\eta}_{\cdot}$ and diagonal covariance matrix (D) . The overall distribution of Y_{ij} , where η_{ij} are treated as random variables, is called a predictive distribution. According to the above assumption, the predictive distribution of each $Y_{i\cdot}$ is normal with the same mean, $\bar{\eta}_{\cdot}$, and covariance matrix $\frac{1}{N} + (D)$ (see Zacks [6; p. 296]). The predictive marginal distributions of the components Y_{ij} are normal with means $\bar{\eta}_j (j=1, \dots, k)$ and variances

$\tau_{ij}^2 = \frac{1}{M_{ij}} + D_j^2 (i=1, \dots, N, j=1, \dots, k)$. Since we assumed that the correlations $\rho_{jj'}$ are the same for all evaluations, we will treat the vectors $\bar{\eta}_{\cdot}$, and the matrices (D) and $(R) = (\rho_{jj'}, j, j' = 1, \dots, k)$ as unknown but fixed parameters. We remark that the predictive marginal distributions of Y_{ij} resemble the distributions of the observations in Model II of Analysis of Variance (see Zacks [6; p. 79]), in which the components of variance are $1/M_{ij}$ (known) and D_j (unknown). We conclude this section by specifying the posterior distributions of η_{ij} given Y_{ij} . According to the Bayesian model, the posterior distribution of η_{ij} given Y_{ij} is normal with mean

$$E \{ \eta_{ij} | Y_{ij} \} = (\bar{\eta}_j + D_j M_{ij} Y_{ij}) / (1 + D_j M_{ij}), \quad (3.1)$$

and variance

$$V \{ \eta_{ij} | Y_{ij} \} = D_j / (1 + M_{ij} D_j) \quad (3.2)$$

Notice that the Bayes estimator of η_{ij} for a squared-error loss is given by (3.1), with a Bayes risk given by (3.2). If $\bar{\eta}_j$ and D_j are unspecified, one can estimate them from the data, to obtain empirical Bayes estimators of the characterizing parameters (see Zacks [6; p. 321]).

4. Bayesian Confidence Intervals for the Characterizing Parameters

Bayesian confidence intervals, at level $1-\alpha$, for specified parameters, are defined as the intervals in the parameterspace whose posterior coverage probabilities, given the sample data, are at least $1-\alpha$. According to the model of Section 3, the posterior coverage probability of the intervals

$$\frac{\bar{\eta}_j + D_j M_{ij} Y_{ij}}{1 + I_j M_{ij}} \pm Z_{1-\alpha/2} \left[\frac{D_j}{1 + M_{ij} D_j} \right]^{1/2} \quad (4.1)$$

is $(1-\alpha)$, where Z_γ is the γ -th fractile of the standard normal distribution. We determine for each category empirical estimates of the prior parameters $\bar{\eta}_j$ and D_j and, by substituting these estimates in (4.1), we obtain empirical Bayes confidence intervals for η_{ij} ($i=1, \dots, N$). The substitution of estimates of $\bar{\eta}_j$ and D_j for the correct values will generally effect the coverage probabilities. The estimates described below are consistent, and we expect that, for large values of N , the actual coverage will be close to the nominal one. In order to provide conservative intervals, we have used in each category, the Bonferroni method of simultaneous confidence intervals, for all the $N = 27$ evaluations (see Miller [4]). For this purpose we replaced α in (4.1) by $\alpha' = \alpha/N$.

Applying the transformation

$$\theta = \sin^2(\eta/2), \quad (4.2)$$

which is strictly increasing for $0 \leq \eta \leq \pi$, we can obtain from the η -confidence intervals θ -confidence intervals. These intervals, for the ten categories and twenty seven evaluations with $\alpha = .05$, are presented in Table 4.1.

Table 4.1

CONFIDENCE INTERVALS FOR CAT. C1-C5

EVAL	1		2		3		4		5	
	L	U	L	U	L	U	L	U	L	U
1	0.802	0.987	0.648	0.942	0.849	0.999	0.366	0.882	0.661	0.926
2	0.902	1.000	0.794	0.982	0.802	0.988	0.481	0.942	0.809	0.988
3	0.834	1.000	0.519	0.942	0.751	0.984	0.333	0.872	0.655	0.940
4	0.853	0.999	0.723	0.991	0.683	0.956	0.556	0.972	0.834	0.999
5	0.548	0.881	0.592	0.931	0.782	0.987	0.598	0.984	0.832	0.999
6	0.814	0.995	0.820	1.000	0.625	0.931	0.699	1.000	0.769	0.980
7	0.791	0.995	0.717	0.980	0.772	0.986	0.437	0.935	0.880	1.000
8	0.397	0.749	0.269	0.668	0.723	0.972	0.318	0.855	0.622	0.912
9	0.778	0.977	0.680	0.964	0.807	0.993	0.717	1.000	0.913	1.000
10	0.792	0.983	0.711	0.968	0.763	0.981	0.517	0.958	0.763	0.975
11	0.755	1.000	0.509	1.000	0.794	0.997	0.714	1.000	0.902	1.000
12	0.634	0.997	0.407	0.982	0.679	0.973	0.688	1.000	0.514	0.858
13	0.716	1.000	0.659	1.000	0.668	0.986	0.328	0.969	0.621	1.000
14	0.900	1.000	0.685	0.984	0.704	0.969	0.427	0.909	0.843	0.999
15	0.834	1.000	0.669	0.996	0.658	0.966	0.432	0.922	0.582	1.000
16	0.768	0.985	0.734	0.992	0.743	0.979	0.335	0.860	0.906	1.000
17	0.791	0.995	0.767	1.000	0.698	0.975	0.432	0.922	0.669	1.000
18	0.920	1.000	0.817	1.000	0.804	0.993	0.682	0.995	0.878	1.000
19	0.901	1.000	0.673	0.957	0.844	0.999	0.793	1.000	0.881	1.000
20	0.747	0.979	0.626	0.972	0.712	0.970	0.321	0.843	0.824	0.996
21	0.586	0.965	0.331	0.929	0.621	0.943	0.128	0.711	0.589	0.903
22	0.871	0.999	0.762	0.997	0.712	0.965	0.568	0.973	0.649	0.924
23	0.871	0.999	0.865	1.000	0.757	0.980	0.682	0.995	0.880	1.000
24	0.823	0.998	0.772	1.000	0.758	0.985	0.609	0.985	0.839	0.999
25	0.770	0.990	0.747	0.997	0.743	0.979	0.396	0.890	0.401	0.754
26	0.753	0.994	0.699	0.991	0.724	0.977	0.634	0.994	0.754	0.979
27	0.921	1.000	0.901	1.000	0.709	0.960	0.691	0.995	0.866	0.999

CONFIDENCE INTERVALS FOR CAT. C6-C10

1	0.807	0.971	0.642	0.921	0.586	0.876	0.771	0.942	0.611	0.962
2	0.872	0.988	0.752	0.966	0.881	0.997	0.789	0.950	0.657	0.973
3	0.675	0.966	0.665	0.946	0.710	0.967	0.804	0.965	0.644	0.979
4	0.844	0.996	0.747	0.979	0.821	0.991	0.808	0.966	0.736	0.997
5	0.840	0.987	0.727	0.971	0.545	0.857	0.793	0.958	0.644	0.979
6	0.827	0.996	0.835	0.997	0.791	0.985	0.819	0.971	0.749	0.997
7	0.877	0.997	0.759	0.984	0.755	0.982	0.825	0.970	0.611	0.962
8	0.385	0.704	0.465	0.812	0.616	0.880	0.759	0.944	0.549	0.940
9	0.746	0.954	0.615	0.914	0.783	0.963	0.797	0.961	0.749	0.997
10	0.723	0.924	0.588	0.892	0.736	0.957	0.789	0.952	0.743	0.997
11	0.733	1.000	0.765	0.999	0.737	1.000	0.811	0.970	0.662	0.987
12	0.451	0.950	0.406	0.843	0.470	0.945	0.788	0.960	0.586	0.968
13	0.671	1.000	0.523	0.986	0.801	1.000	0.794	0.966	0.640	0.993
14	0.744	0.974	0.700	0.957	0.746	0.967	0.805	0.965	0.594	0.952
15	0.856	1.000	0.747	0.994	0.775	0.993	0.811	0.970	0.580	0.951
16	0.894	1.000	0.796	0.987	0.763	0.974	0.819	0.971	0.604	0.961
17	0.895	1.000	0.823	0.999	0.856	1.000	0.808	0.966	0.714	0.996
18	0.876	0.999	0.850	0.997	0.869	0.996	0.826	0.973	0.658	0.980
19	0.812	0.978	0.701	0.944	0.737	0.937	0.809	0.966	0.743	0.997
20	0.654	0.934	0.532	0.863	0.689	0.944	0.797	0.961	0.705	0.989
21	0.464	0.928	0.463	0.863	0.323	0.797	0.792	0.961	0.549	0.940
22	0.815	0.982	0.670	0.947	0.837	0.989	0.829	0.974	0.717	0.990
23	0.909	1.000	0.789	0.983	0.862	0.993	0.804	0.963	0.691	0.989
24	0.851	0.999	0.758	0.980	0.688	0.954	0.793	0.961	0.698	0.989
25	0.733	0.968	0.704	0.958	0.653	0.937	0.804	0.965	0.672	0.981
26	0.704	0.964	0.617	0.943	0.617	0.923	0.814	0.970	0.684	0.988
27	0.933	1.000	0.847	0.994	0.924	0.999	0.810	0.964	0.711	0.990

The empirical estimates for $\bar{\eta}_j$ and D_j applied here are:

$$\hat{\eta}_{j,N} = \frac{1}{N} \sum_{i=1}^N Y_{ij}^2, \quad (4.3)$$

and

$$\hat{D}_{j,N} = \left[\frac{1}{N} \sum_{i=1}^N Y_{ij}^2 - \hat{\eta}_{j,N}^2 - H_{j,N}^{-1} \right]^+, \quad (4.4)$$

where $[a]^+ = \max(a, 0)$, and $H_{j,N}$ is the harmonic mean of M_{ij} , i.e., $H_{j,N}^{-1} = \frac{1}{N} \sum_{i=1}^N \frac{1}{M_{ij}}$. Notice that $\hat{\eta}_{j,N}$ is a strongly consistent estimator of $\bar{\eta}_j$ (approaches $\bar{\eta}_j$ a.s. as $N \rightarrow \infty$). Assuming that $H_{j,N}$ converges to a limit as $N \rightarrow \infty$, $\hat{D}_{j,N}$ converges to D_j as N grows. The values of $\hat{\eta}_{j,N}$ and $D_{j,N}$ obtained in this manner are given in Table 4.2.

Table 4.2
Empirical Estimates
of $\bar{\eta}_j$ and D_j

j	$\bar{\eta}_{j,N}$	$\hat{D}_{j,N}$
1	2.5851	0.0755
2	2.4125	0.1058
3	2.4494	0.0297
4	2.1691	0.1464
5	2.5379	0.0871
6	2.5185	0.0994
7	2.3468	0.0751
8	2.4176	0.0917
9	2.4886	0.0086
10	2.3883	0.0388

5. Estimating the Correlations

The marginal bivariate predictive distribution of Y_{ij} and $Y_{ij'}$, for $j \neq j'$ is normal with p.d.f.

$$f(Y_{ij}, Y_{ij'}; \bar{\eta}_j, \bar{\eta}_{j'}, \rho_{jj'}, D_j, D_{j'}) =$$

$$\frac{1}{2\pi} \cdot \left(\frac{M_{ij}}{1 + D_j M_{ij}} \right)^{1/2} \left(\frac{M_{ij'}}{1 + D_{j'} M_{ij'}} \right)^{1/2} \cdot \frac{1}{\sqrt{1 - \rho_{jj'}^2}} \cdot$$

$$\exp \left\{ - \frac{1}{2(1 - \rho_{jj'}^2)} \left[\frac{M_{ij}}{1 + M_{ij} D_j} (Y_{ij} - \bar{\eta}_j)^2 + \frac{M_{ij'}}{1 + M_{ij'} D_{j'}} (Y_{ij'} - \bar{\eta}_{j'})^2 \right. \right.$$

$$\left. \left. - 2 \rho_{jj'} \frac{(Y_{ij} - \bar{\eta}_j) \sqrt{M_{ij}}}{\sqrt{1 + M_{ij} D_j}} \cdot \frac{(Y_{ij'} - \bar{\eta}_{j'}) \sqrt{M_{ij'}}}{\sqrt{1 + M_{ij'} D_{j'}}} \right] \right\} \quad (5.1)$$

Define the weight functions

$$w_{ij} = \frac{M_{ij}}{1 + M_{ij} D_j}, \quad i=1, \dots, N, \quad j=1, \dots, K. \quad (5.2)$$

If the prior means and variances, $\bar{\eta}_j, D_j$ ($j=1, \dots, k$) are known, then, the predictive quasi-likelihood function of $\rho_{jj'}$, given the sample data is

$$L(\rho_{jj'} | Y_{ij}, Y_{ij'}, i=1, \dots, N) =$$

$$(1 - \rho_{jj'}^2)^{N/2} \exp \left\{ - \frac{1}{2(1 - \rho_{jj'}^2)} \left[\sum_{i=1}^N w_{ij} (Y_{ij} - \bar{\eta}_j)^2 + \right. \right.$$

$$\left. \left. \sum_{i=1}^N w_{ij'} (Y_{ij'} - \bar{\eta}_{j'})^2 - 2 \rho_{jj'} \sum_{i=1}^N \sqrt{w_{ij} w_{ij'}} (Y_{ij} - \bar{\eta}_j) (Y_{ij'} - \bar{\eta}_{j'}) \right] \right\} \quad (5.3)$$

(5.3) is called a 'quasi-likelihood' since the correct likelihood function is based on the k -variate distributions of $Y_{\cdot 1}, \dots, Y_{\cdot N}$.

In practice, $\bar{\eta}_j$ and D_j are unknown, we will therefore use the quasi-likelihood function of ρ_{jj} , by substituting in (5.3) estimates of $\bar{\eta}_j$ and D_j . Thus, let

$$S_j^2 = \sum_{i=1}^N \hat{W}_{ij} (Y_{ij} - \hat{\eta}_j)^2, \quad j=1, \dots, k$$

and

(5.4)

$$P_{jj'} = \sum_{i=1}^N \sqrt{\hat{W}_{ij} \hat{W}_{ij'}} (Y_{ij} - \hat{\eta}_j) (Y_{ij'} - \hat{\eta}_{j'}), \quad j \neq j'$$

where

$$\hat{\eta}_j = \frac{\sum_{i=1}^N \hat{W}_{ij} Y_{ij}}{\sum_{i=1}^N \hat{W}_{ij}}, \quad (5.5)$$

and

$$\hat{W}_{ij} = M_{ij} / (1 + M_{ij} \hat{D}_{j,N}) \quad (5.6)$$

The estimator $\hat{\rho}_{j,j'}$, which maximizes the log-quasi-likelihood

$$\ell(\rho_{jj'}) = \frac{N}{2} \log (1 - \rho_{jj'}^2) - \frac{1}{2(1 - \rho_{jj'}^2)} \left[S_j^2 + S_{j'}^2 - 2\rho_{jj'} P_{jj'} \right], \quad (5.7)$$

is called quasi-maximum likelihood (q.m.l.) estimator of $\rho_{jj'}$. The derivative of $\ell(\rho)$ with respect to ρ is

$$\frac{\partial \ell(\rho)}{\partial \rho} = N \frac{\rho}{1 - \rho^2} - \frac{\rho(S_j^2 + S_{j'}^2) - P_{jj'}(1 + \rho^2)}{(1 - \rho^2)^2}. \quad (5.8)$$

Accordingly, the q.m.l. estimator of $\rho_{jj'}$ is a solution of the cubic equation

$$\rho^3 - \frac{P_{jj'}}{N} \rho^2 + \left(\frac{S_j^2}{n} + \frac{S_{j'}^2}{n} - 1 \right) \rho - \frac{P_{jj'}}{n} = 0 \quad (5.9)$$

for which the second derivative of (5.7) is negative. Accordingly, for given values of $\hat{D}_{j,N}$ ($j=1, \dots, k$), we determine \hat{W}_{ij} ($i=1, \dots, N_j$ $j=1, \dots, k$), and $\hat{\eta}_j$ ($j=1, \dots, k$) and solve (5.9) by the Newton-Raphson method, starting with the initial solutions

$$\tilde{\rho}_j = 2P_{jj'} / (S_j^2 + S_{j'}^2), \quad j=1, \dots, k \quad (5.10)$$

In Table 5.1 we provide the initial estimates of the correlations (5.10) and the final estimates, which are the solutions of (5.9). The estimates of D_j are those given in Table 4.2. Two numbers are given in each cell of the matrix of final estimates of the correlations. The upper number is the correlation and the lower number is an estimate of its standard error, namely

$$S.E. \{ \hat{\rho}_{jj'} \} = (1 - \hat{\rho}_{jj'}^2) / \sqrt{N(1 + \hat{\rho}_{jj'}^2)} \quad (5.11)$$

Every estimate $\hat{\rho}_{jj'}$, which, in absolute value, is greater than $2 S.E. \{ \hat{\rho}_{jj'} \}$ is significant.

Table 5.1

Initial and final estimates of the
correlation parameters

INITIAL ESTIMATES OF CORRELATIONS

1.000	0.765	0.168	0.360	0.373	0.678	0.593	0.692	0.616	0.406
0.765	1.000	0.029	0.442	0.444	0.851	0.781	0.763	0.635	0.593
0.168	0.029	1.000	0.321	0.321	0.183	0.122	0.099	-0.133	0.140
0.360	0.442	0.321	1.000	0.385	0.341	0.336	0.381	0.305	0.565
0.373	0.444	0.321	0.385	1.000	0.613	0.560	0.542	0.364	0.354
0.678	0.851	0.183	0.341	0.613	1.000	0.924	0.721	0.623	0.386
0.593	0.781	0.122	0.336	0.560	0.924	1.000	0.719	0.641	0.333
0.692	0.763	0.099	0.381	0.542	0.721	0.719	1.000	0.491	0.489
0.616	0.635	-0.133	0.305	0.364	0.623	0.641	0.491	1.000	0.322
0.406	0.593	0.140	0.565	0.354	0.386	0.333	0.489	0.322	1.000

FINAL ESTIMATES OF CORRELATIONS

1	1.000	0.760	0.163	0.355	0.348	0.676	0.587	0.694	0.583	0.394
	0.0	0.065	0.185	0.158	0.160	0.087	0.109	0.082	0.110	0.151
2	0.760	1.000	0.029	0.443	0.421	0.853	0.781	0.769	0.607	0.587
	0.065	0.0	0.192	0.141	0.146	0.040	0.059	0.062	0.104	0.109
3	0.163	0.029	1.000	0.319	0.299	0.183	0.120	0.101	-0.120	0.136
	0.185	0.192	0.0	0.165	0.168	0.183	0.188	0.190	0.188	0.187
4	0.355	0.443	0.319	1.000	0.366	0.347	0.339	0.392	0.283	0.561
	0.158	0.141	0.165	0.0	0.157	0.160	0.161	0.152	0.170	0.115
5	0.348	0.421	0.299	0.366	1.000	0.596	0.538	0.527	0.322	0.330
	0.160	0.146	0.168	0.157	0.0	0.107	0.120	0.123	0.164	0.163
6	0.676	0.853	0.183	0.347	0.596	1.000	0.926	0.732	0.599	0.385
	0.087	0.040	0.183	0.160	0.107	0.0	0.020	0.072	0.106	0.153
7	0.587	0.781	0.120	0.339	0.538	0.926	1.000	0.726	0.615	0.328
	0.109	0.059	0.188	0.161	0.120	0.020	0.0	0.074	0.102	0.163
8	0.694	0.769	0.101	0.392	0.527	0.732	0.726	1.000	0.469	0.492
	0.082	0.062	0.190	0.152	0.123	0.072	0.074	0.0	0.136	0.131
9	0.583	0.607	-0.120	0.283	0.322	0.599	0.615	0.469	1.000	0.294
	0.110	0.104	0.188	0.170	0.164	0.106	0.102	0.136	0.0	0.169
10	0.394	0.587	0.136	0.561	0.330	0.385	0.328	0.492	0.294	1.000
	0.151	0.109	0.187	0.115	0.163	0.153	0.163	0.131	0.169	0.0

6. Summary

In the current paper we present confidence intervals for the category scores as well as the correlations between the scores. The category scores help determine the weaknesses and the strengths of the units. It is important to obtain confidence intervals on these scores because before planning remedial training it is necessary to know whether or not the scores are meaningful. Thus high correlation between two categories may indicate that training to improve the performance in one can improve the performance in the other.

The results in Table 5.1 are intuitively explainable. We obtain high correlation between the categories evaluating the command's performance - Categories 1, 2, 6, 7, and 8. But Categories 3 (Communicating), 4 (Performing as Marines), 5 (Delivering supporting fire) and 10 (Supervising required actions of individual Marines) evaluate the performance of various specific groups and thus each of them cannot be expected to have high correlation with other categories.

A major finding - see Appendix A - is that, for any one battalion, the probability of "Yes" for a given requirement is the same for applicable requirements of the same category. This provides considerable justification for the use of 10 category scores as a measure of the overall capability of the battalion.

APPENDIX 1

We consider here the question of testing assumption A.2.

Typically, for a given evaluation, there are a certain number of "Yes's" among the applicable requirements. The question is how to test the hypothesis that the probability of a "Yes" in a given evaluation depends only on the category to which the requirement belongs. We have tested whether this hypothesis is supported by the data, in the following manner.

The data sets corresponding to the $N = 27$ different evaluations are arranged according to tasks within 17 different Military Performance Standards (MPS; see [7]). We therefore considered first the number of "Yes's" among the applicable requirements within each cell of MPS x category, for each evaluation. In Table A.1 we present these statistics, as an example, for EVAL 1. An exact test of the hypothesis that the probability of "Yes" is the same for all the MPS's within a category would be very difficult to perform, due to the small number of applicable requirements in each cell.

Let θ_{ikj} denote the probability of "Yes" in EVAL i , MPS k , Category j ($i=1, \dots, 27$; $k=1, \dots, 17$; $j=1, \dots, 10$). We consider the set of null hypotheses:

$$H_0^{(i,j)}: \theta_{ikj} = \theta_{ij} \text{ for all } k=1, \dots, 17 ;$$

$$\text{at each } i=1, \dots, 27 ; j=1, \dots, 10 .$$

In the spirit of the Bayesian approach of Section 3, we assume, a priori, that θ_{ikj} are independent random variables, having some prior distributions

$H_{kj}(\theta)$ ($k=1, \dots, 17$; $j=1, \dots, 10$). If the null hypotheses $H_0^{(i,j)}$ are true for all (i,j) , the following null hypotheses should also be true: For each j , $j=1, \dots, 10$,

$$H_0^{(\cdot,j)}: \psi_{kj} = \psi_j \text{ for all } k=1, \dots, 17 ,$$

where

$$\psi_{kj} = \int_0^1 \theta_{ijk} dH_{kj}^{(\theta)}(\theta_{ijk}) \quad (A.1.1)$$

Table A.1

The number of satisfied (upper) and applicable (lower)
requirement by MPS and Category for Eval. 1

MPS	CATEGORIES									
	1	2	3	4	5	6	7	8	9	10
2.A.1	9	4	4	11	0	3	14	4	2	6
	11	9	4	19	0	3	18	13	3	10
2.A.2	8	0	20	0	7	10	15	9	9	3
	9	0	20	1	8	10	18	9	9	3
2.A.3	0	0	2	0	35	3	0	0	0	0
	0	0	2	0	48	5	1	0	0	0
2.B.1	6	0	0	0	0	5	2	5	0	2
	6	0	0	0	0	5	3	5	1	2
2.B.2	7	5	1	2	0	8	3	7	2	1
	7	6	1	2	0	9	3	10	2	1
2.B.3	8	5	3	0	1	2	1	11	0	0
	8	5	3	0	1	3	1	14	0	0
2.B.4	9	3	1	0	2	9	5	12	2	1
	10	5	1	0	2	11	8	13	2	1
2.B.5	0	0	0	0	0	0	0	0	0	0
	0	0	0	0	0	0	0	0	0	0
2.B.6	2	3	2	1	2	10	2	0	0	0
	2	4	2	1	2	12	3	0	0	0
2.B.7	3	2	0	0	0	6	0	1	2	0
	4	2	0	0	0	7	0	2	3	0
2.C.1	0	0	0	0	0	0	0	0	0	0
	0	0	0	0	0	0	0	0	0	0
2.C.2	3	3	1	0	0	4	1	1	0	0
	3	3	1	0	0	4	1	4	0	0
2.C.3	0	0	0	0	0	0	0	0	0	0
	0	0	0	0	0	0	0	0	0	0
2.C.4	3	2	0	0	0	3	4	0	0	0
	3	2	0	0	0	3	4	0	0	0
2.D.1	2	3	6	0	8	34	0	0	4	0
	2	3	6	0	8	34	0	0	4	0
2.D.2	2	6	0	0	0	2	3	3	3	0
	2	6	0	0	0	2	3	3	8	0
2.D.3	0	8	0	0	0	1	0	0	0	0
	0	9	0	0	0	2	0	0	0	0
VOL. II	62	44	40	14	55	100	50	53	24	13
	67	54	40	23	69	110	63	73	32	17

Let ψ_{kj}^* and M_{kj}^* ($k=1, \dots, 17$, $j=1, \dots, 10$) be the number of satisfied and applicable requirements in the k -th MPS and j -th category, summed over all the 27 evaluations. Let

$$X_{\cdot j}^* = \sum_{k=1}^{17} X_{kj}^* \quad (j=1, \dots, 10) \quad (A.1.2)$$

and

$$M_{\cdot j}^* = \sum_{k=1}^{17} M_{kj}^*$$

These statistics are given in Table A.2. We see in Table A.2 that the values of $M_{\cdot j}^*$ are large ones, for all $j=1, \dots, 10$; and that M_{kj}^* are generally not too small. We therefore consider, for each hypothesis $H_0^{(\cdot, j)}$, the chi-squared statistic

$$T_j^2 = \sum_{k=1}^{k_j^*} \frac{(X_{kj}^* - E_{kj}^*)^2}{E_{kj}^*} I_{kj}^* \quad , \quad (A.1.3)$$

where

$$E_{kj}^* = M_{kj}^* \frac{X_{\cdot j}^*}{M_{\cdot j}^*} \quad , \quad \begin{matrix} k=1, \dots, 17 \\ j=1, \dots, 10 \end{matrix} \quad (A.1.4)$$

$$I_{kj}^* = \begin{cases} 1 & , \text{ if } M_{kj}^* > 0 \\ 0 & , \text{ if } M_{kj}^* = 0 \end{cases} \quad (A.1.5)$$

and $k_j^* = \sum_{k=1}^{17} I_{kj}^*$. The estimates of the expected values of X_{kj}^* , under the null hypothesis $H_0^{(\cdot, j)}$, are E_{kj}^* . The large sample distribution of T_j^2 is like that of a chi-squared random variable with $v_j = k_j^* - 1$ degrees of freedom. In Table A.3 we provide the values of T_j^2 and their P -values

$$P_j = P \{X^2[v_j] \geq T_j^2 | T_j^2\} \quad , \quad j=1, \dots, 10 \quad (A.1.6)$$

Table A.2

The statistics $X_{.j}^*$ and $M_{.j}^*$

1	214	158	66	366	0	65	351	216	64	219
1	245	199	93	472	0	70	413	283	75	258
2	212	0	416	18	162	218	367	204	207	70
2	232	0	447	26	182	239	440	228	225	75
3	0	0	36	0	993	77	21	0	0	0
3	0	0	49	0	1121	106	24	0	0	0
4	63	0	0	7	13	48	24	74	24	18
4	63	0	0	9	13	49	27	77	27	18
5	110	99	11	33	24	142	51	162	20	26
5	121	110	15	37	25	164	54	189	22	34
6	101	64	39	0	10	32	9	180	0	0
6	113	74	40	0	10	37	10	193	0	0
7	225	123	22	0	36	260	178	347	43	20
7	243	140	22	0	40	282	211	372	43	21
8	48	54	30	22	0	97	62	70	7	0
8	53	68	31	26	0	105	72	89	8	0
9	118	89	40	38	30	261	96	86	31	0
9	123	101	44	51	33	293	108	94	32	0
10	81	26	0	0	5	132	0	80	32	0
10	83	26	0	0	5	142	0	95	37	0
11	48	31	11	0	2	74	0	104	0	19
11	48	33	12	0	2	76	0	107	0	21
12	49	87	19	0	13	75	30	59	0	0
12	54	100	19	0	15	90	36	75	0	0
13	21	17	7	0	9	30	4	41	5	0
13	24	18	7	0	10	36	4	44	6	0
14	28	22	0	0	0	37	26	0	0	0
14	30	23	0	0	0	38	32	0	0	0
15	10	19	29	0	40	178	0	0	20	0
15	10	19	30	0	40	181	0	0	20	0
16	8	24	2	0	0	5	8	12	20	0
16	8	24	2	0	0	5	10	12	28	0
17	0	39	3	0	0	11	0	0	0	0
17	0	43	3	0	0	13	0	0	0	0
TOTAL I	1336	852	731	484	1337	1742	1227	1635	473	372
TOTAL N	1450	978	814	621	1496	1926	1441	1858	523	427

where $\chi^2[v_j]$ is a chi-squared random variable, independent of T_j^2 , with v_j degrees of freedom

Table A.3

The values of T_j^2 and P_j

Cat.	T_j^2	P_j	Cat.	T_j^2	P_j
1	2.155	.99987	6	7.371	.96543
2	3.811	.99304	7	1.200	.99996
3	7.326	.88458	8	10.183	.59991
4	1.089	.95507	9	2.292	.99357
5	1.112	.99991	10	1.452	.91854

We see in Table A.3 that each one of the ten null-hypotheses $H_0^{(\cdot,j)}$ is strongly accepted (very high P - values). Let P^* be the combined P - value of all the ten tests. According to the Bonferroni inequality [4],

$$P^* \geq 1 - \sum_{j=1}^{10} (1-P_j) \quad . \quad (A.1.7)$$

Thus, according to the values in Table A.3, $P^* \geq .30988$. Thus, we simultaneously accept all the ten null hypotheses, with a very high P^* value. Note that, if at least one of the null hypotheses had been rejected we should have concluded that assumption A.2 of Section 2 is not supported by the data. However, since all the null hypotheses have been strongly accepted, we conclude that the data is in accord with assumption A.2.

APPENDIX 2

We show in the present appendix that asymptotically, the correlation between $\hat{\theta}_{ij}$ and $\hat{\theta}_{ij'}$, is equivalent to $\rho_{jj'}$, for $j, j'=1, \dots, k$.

The inverse of (2.1) is

$$\hat{\theta}_{ij} = \sin^2(Y_{ij}/2), \quad \begin{matrix} i=1, \dots, N \\ j=1, \dots, k \end{matrix} \quad (\text{A.2.1})$$

where $Y_{ij} \sim N(\eta_{ij}, M_{ij}^{-1})$. Expand $\hat{\theta}_{ij}$ around η_{ij} to obtain

$$\begin{aligned} \hat{\theta}_{ij} &= \sin^2(\eta_{ij}/2) + (Y_{ij} - \eta_{ij}) \sin(\eta_{ij}/2) \cos(\eta_{ij}/2) \\ &\quad + O_P\left(\frac{1}{\sqrt{N}}\right), \end{aligned} \quad (\text{A.2.2})$$

where $O_P(\cdot)$ is an order-of-magnitude-in-probability symbol (see Zacks [5; pp. 208]).

Let $\hat{\theta}_{\sim i} = (\hat{\theta}_{i1}, \dots, \hat{\theta}_{ik})'$ and $\hat{\tau}_{\sim i} = \left(\sin^2\left(\frac{\eta_{i1}}{2}\right), \dots, \sin^2\left(\frac{\eta_{ik}}{2}\right)\right)'$.

Notice that $\sin\left(\frac{\eta}{2}\right) \cos\left(\frac{\eta}{2}\right) = \frac{1}{2} \sin(\eta)$. Let $(D)_i =$

$\text{diag}\{\sin(\eta_{i1}), \dots, \sin(\eta_{ik})\}$.

Then,

$$\hat{\theta}_{\sim i} = \hat{\tau}_{\sim i} + \frac{1}{2} (D)_i (Y_{\sim i} - \eta_{\sim i}) + O_P\left(\frac{1}{\sqrt{N}}\right), \quad (\text{A.2.3})$$

Thus, as $N \rightarrow \infty$, the asymptotic covariance matrix of $\hat{\theta}_{\sim i}$ is

$$\ddagger(\hat{\theta}_{\sim i}) = \frac{1}{4} (D)_i \ddagger_1 (D)_i, \quad (\text{A.2.4})$$

where the elements of \ddagger_1 are specified in (2.2). Accordingly, the asymptotic variance of $\hat{\theta}_{ij}$ is

$$AV \{\hat{\theta}_{ij}\} = \frac{1}{4} \sin^2(\eta_{ij})/M_{ij} \quad (A.2.5)$$

and the asymptotic covariance of $\hat{\theta}_{ij}$ and $\hat{\theta}_{ij'}$, $j \neq j'$, is

$$AC(\hat{\theta}_{ij}, \hat{\theta}_{ij'}) = \frac{1}{4} \rho_{jj'} \sin(\eta_{ij}) \sin(\eta_{ij'}) / \sqrt{M_{ij} M_{ij'}} \quad (A.2.6)$$

Hence, the asymptotic correlation between $\hat{\theta}_{ij}$ and $\hat{\theta}_{ij'}$ is $\rho_{jj'}$, for all $i=1, \dots, N$.

REFERENCES

- [1] Barzily, Z. (1980). Analyzing MCCRES data. Technical Paper Serial T-427.
- [2] Barzily, Z., P. R. Catalogne, and W. H. Marlow (1981). Assessing Marine Corps Readiness. Defense Management Journal, Vol. 17, No. 1, pp. 25-29.
- [3] Johnson, N. L and S. I. Kotz (1970). Discrete Distributions: Distributions in Statistics. Wiley.
- [4] Miller Jr., R. G. (1981). Simultaneous Statistical Inference. 2nd Ed., Springer.
- [5] Zacks, S. (1971). The Theory of Statistical Inference. Wiley.
- [6] Zacks, S. (1981). Parametric Statistical Inference: Basic Theory and Modern Approaches. Pergamon.
- [7] Zacks, S. and W. H. Marlow (1981). Statistical problems in analyzing Marine Corps readiness evaluations. Technical Memorandum Serial TM-68526.
- [8] Zacks, S., W. H. Marlow, and Z. Barzily (1981). Factor analysis of task-scores in Marine Corps readiness evaluations. Technical Memorandum Serial TM-69195.

THE GEORGE WASHINGTON UNIVERSITY
Program in Logistics
Distribution List for Technical Papers

The George Washington University Office of Sponsored Research Gelman Library Vice President H. F. Bright Dean Harold Liebowitz Dean Henry Solomon	Armed Forces Industrial College	Case Western Reserve University Prof B. V. Dean Prof M. Mesarovic
ONR Chief of Naval Research (Codes 200, 434) Resident Representative	Armed Forces Staff College	Cornell University Prof R. E. Beehoffer Prof R. W. Conway Prof Andrew Schultz, Jr.
OPNAV OP-40 DCNO, Logistics Navy Dept Library NAVDATA Automation Cmd	Army War College Library Carlisle Barracks	Cowles Foundation for Research in Economics Prof Martin Shubik
Naval Aviation Integrated Log Suppo	Army Cmd & Gen Staff College	Florida State University Prof K. A. Bradley
NARDAC Tech Library	Army Logistics Mgt Center Fort Lee	Harvard University Prof W. G. Cochran Prof Arthur Schleifer, Jr.
Naval Electronics Lab Library	Commanding Officer, USALDSRA New Cumberland Army Depot	Princeton University Prof A. W. Tucker Prof J. W. Tukey Prof Geoffrey S. Watson
Naval Facilities Eng Cmd Tech Library	Army Inventory Res Ofc Philadelphia	Purdue University Prof S. S. Gupta Prof H. Rubin Prof Andrew Whinston
Naval Ordnance Station Louisville, Ky. Indian Head, Md.	Army Trans Material Cmd TCMAC-ASDT	Stanford University Prof T. W. Anderson Prof Kenneth Arrow Prof G. B. Dantzig Prof F. S. Hillier Prof D. L. Ichter Prof Samuel Lin Prof G. J. L. Lerman Prof Herbert Solomon Prof A. E. Veinott, Jr.
Naval Ordnance Sys Cmd Library	Air Force Headquarters AFADS-3 LEXY SAF/ALG	University of California, Berkeley Prof R. E. Barlow Prof D. Gale Prof Jack Kiefer
Naval Research Branch Office Boston Chicago New York Pasadena San Francisco	Griffiss Air Force Base Reliability Analysis Center	University of California, Los Angeles Prof R. E. O'Neill
Naval Ship Eng Center Philadelphia, Pa.	Gunter Air Force Base AFLMC/XR	University of North Carolina Prof W. L. Smith Prof M. K. Leadbetter
Naval Ship Res & Dev Center	Maxwell Air Force Base Library	University of Pennsylvania Prof Russell Ackoff
Naval Sea Systems Command PMS 30611 Tech Library Code 073	Wright-Patterson Air Force Base AFLC/OA Research Sch Log AFALD/XR	University of Texas Institute for Computing Science and Computer Applications
Naval Supply Systems Command Library Operations and Inventory Analysis	Defense Technical Info Center	Yale University Prof E. L. Anscombe Prof H. Scott
Naval War College Library Newport	National Academy of Sciences Maritime Transportation Res Ctr	Prof Z. W. Birnbaum University of Washington
BIPERS Tech Library	National Bureau of Standards Dr B. H. Colvin Dr Joan Rosenblatt	Prof B. H. Bissinger The Pennsylvania State University
FMSO	National Science Foundation	Prof Seth Bonder University of Michigan
USN Ammo Depot Earle	National Security Agency	Prof G. E. Box University of Wisconsin
USN Postgrad School Monterey Library Dr Jack R. Borsting Prof C. R. Jones	Weapons Systems Evaluation Group	Dr Jerome Bracken Institute for Defense Analyses
US Coast Guard Academy Capt Jimmie D. Woods	British Navy Staff	
US Marine Corps Commandant Deputy Chief of Staff, R&D	National Defense Hdqtrs, Ottawa Logistics, OR Analysis Etab	
Marine Corps School Quantico Landing Force Dev Ctr Logistics Officer	American Power Jet Co George Chernowitz	
	General Dynamics, Pomona	
	General Research Corp Library	
	Logistics Management Institute Dr Murray A. Weisler	
	Rand Corporation Library Mr William F. Butzler	
	Carnegie-Mellon University Dean R. A. Simon Prof G. Thompson	

Continued

Prof A. Charnes
University of Texas

Prof H. Chernoff
Mass Institute of Technology

Prof Arthur Cohen
Rutgers - The State University

Mr Wallace M. Cohen
US General Accounting Office

Prof C. Derman
Columbia University

Prof Masao Fukushima
Kyoto University

Prof Saul I. Gass
University of Maryland

Dr Donald P. Gaver
Carmel, California

Prof Amrit L. Goel
Syracuse University

Prof J. F. Hannan
Michigan State University

Prof H. O. Hartley
Texas A & M Foundation

Prof W. M. Hirsch
Courant Institute

Dr Alan J. Hoffman
IBM, Yorktown Heights

Prof John R. Isbell
SUNY, Amherst

Dr J. L. Jain
University of Delhi

Prof J. H. K. Kao
Polytech Institute of New York

Prof W. Kruskal
University of Chicago

Mr S. Kumar
University of Madras

Prof C. E. Lemke
Rensselaer Polytech Institute

Prof Loynes
University of Sheffield, England

Prof Tom Maul
Kowloon, Hong Kong

Prof Steven Nahmias
University of Santa Clara

Prof D. B. Owen
Southern Methodist University

Prof P. R. Parathasarathy
Indian Institute of Technology

Prof E. Parzen
Texas A & M University

Prof H. O. Posten
University of Connecticut

Prof R. Remage, Jr.
University of Delaware

Prof Hans Riedwyl
University of Berne

Mr David Rosenblatt
Washington, D. C.

Prof M. Rosenblatt
University of California, San Diego

Prof Alan J. Rowe
University of Southern California

Prof A. H. Rubenstein
Northwestern University

Prof Thomas L. Saaty
University of Pittsburgh

Dr M. E. Salveson
West Los Angeles

Prof Gary Scudder
University of Minnesota

Prof Edward A. Silver
University of Waterloo, Canada

Prof Rosedith Sitgreaves
Washington, DC

LTC G. L. Slyman, MSC
Department of the Army

Prof M. J. Sobel
Georgia Inst of Technology

Prof R. M. Thrall
Rice University

Dr S. Vajda
University of Sussex, England

Prof T. M. Whitin
Wesleyan University

Prof Jacob Wolfowitz
University of South Florida

Prof Max A. Woodbury
Duke University

Prof S. Zacks
SUNY, Binghamton

Dr Israel Zang
Tel-Aviv University

END

FILMED

1-83

DTIC